

A study on the sum of the squares of generalized balancing numbers: the sum formula $\sum_{k=0}^n x^k W_{mk+j}^2$

Yüksel Soykan ¹, Erkan Taşdemir  ² and Can Murat Dikmen ¹

¹Zonguldak Bülent Ecevit University, Department of Mathematics, Art and Science Faculty, 67100, Zonguldak, Turkey

²Kırklareli University, Pınarhisar Vocational School, 39300, Kırklareli, Turkey

Received 16 November 2021, Accepted 24 December 2021, Published 27 December 2021

Abstract. In this paper, closed forms of the sum formulas $\sum_{k=0}^n x^k W_{mk+j}^2$ for generalized balancing numbers are presented. As special cases, we give sum formulas of balancing, modified Lucas-balancing and Lucas-balancing numbers.

Keywords: Balancing numbers, modified Lucas-balancing numbers, Lucas-balancing numbers, sum formulas.

2020 Mathematics Subject Classification: 11B37, 11B39, 11B83.

1 Introduction

A generalized balancing sequence $\{W_n\}_{n \geq 0} = \{W_n(W_0, W_1)\}_{n \geq 0}$ is defined by the second-order recurrence relation

$$W_n = 6W_{n-1} - W_{n-2} \quad (1.1)$$

with the initial values $W_0 = c_0, W_1 = c_1$ not all being zero.

The sequence $\{W_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$W_{-n} = 6W_{-(n-1)} - W_{-(n-2)}$$

for $n = 1, 2, 3, \dots$. Therefore, recurrence (1.1) holds for all integer n .

The Binet formula of generalized balancing numbers can be written as

$$W_n = \frac{W_1 - \beta W_0}{\alpha - \beta} \alpha^n - \frac{W_1 - \alpha W_0}{\alpha - \beta} \beta^n,$$

where α and β are the roots of the quadratic equation $x^2 - 6x + 1 = 0$. Moreover

$$\begin{aligned} \alpha &= 3 + 2\sqrt{2}, \\ \beta &= 3 - 2\sqrt{2}. \end{aligned}$$

 Corresponding author. Email: erkantasdemir@hotmail.com

Note that

$$\begin{aligned}\alpha + \beta &= 6, \\ \alpha\beta &= 1, \\ \alpha - \beta &= 4\sqrt{2}.\end{aligned}$$

Now, one defines three special cases of the sequence $\{W_n\}$. Balancing sequence $\{B_n\}_{n \geq 0}$, modified Lucas-balancing sequence $\{H_n\}_{n \geq 0}$ and Lucas-balancing sequence $\{C_n\}_{n \geq 0}$ are defined, respectively, by the second-order recurrence relations

$$B_n = 6B_{n-1} - B_{n-2}, \quad B_0 = 0, B_1 = 1, \quad (1.2)$$

$$H_n = 6H_{n-1} - H_{n-2}, \quad H_0 = 2, H_1 = 6, \quad (1.3)$$

$$C_n = 6C_{n-1} - C_{n-2}, \quad C_0 = 1, C_1 = 3. \quad (1.4)$$

The sequences $\{B_n\}_{n \geq 0}$, $\{H_n\}_{n \geq 0}$ and $\{C_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$\begin{aligned}B_{-n} &= 6B_{-(n-1)} - B_{-(n-2)}, \\ H_{-n} &= 6H_{-(n-1)} - H_{-(n-2)}, \\ C_{-n} &= 6C_{-(n-1)} - C_{-(n-2)},\end{aligned}$$

for $n = 1, 2, 3, \dots$ respectively. Therefore, recurrences (1.2)-(1.4) hold for all integer n . For more information on generalized balancing numbers, see Soykan [29].

In [1], Behera and Panda defined balancing numbers n as solutions of the diophantine equation

$$1 + 2 + \dots + (n-1) = (n+1) + (n+2) + \dots + (n+r)$$

for some natural number r , called the balancer corresponding to n . The n th balancing number is denoted by B_n . Moreover, $C_n = \sqrt{8B_n^2 + 1}$ is called the n th Lucas-balancing number (see [16]). In fact, B_n and C_n satisfy the second order linear recurrence relations (1.2) and (1.4) respectively. $(B_n)_{n \geq 0}$ is the sequence A001109 in the OEIS [27], whereas $(C_n)_{n \geq 0}$ is the id-number A001541 in OEIS. Balancing and Lucas-balancing sequences have been studied by many authors and more detail can be found in the extensive literature dedicated to these sequences, see for example, [1-4, 9-26].

2 The Sum Formula $\sum_{k=0}^n x^k W_{mk+j}^2$

The following theorem presents sum formulas of generalized balancing numbers.

Theorem 2.1. *Let x be a real (or complex) number. For all integers m and j , for generalized balancing numbers (the case $r = 6$, $s = -1$), the following sum formulas hold:*

(a) *If $(1 + (-s)^{2m}x^2 - xH_{2m})((-s)^m x - 1) \neq 0$, then*

$$\sum_{k=0}^n x^k W_{mk+j}^2 = \frac{\Omega_1}{32(1 + x^2 - xH_{2m})(x - 1)}, \quad (2.1)$$

where

$$\Omega_1 = 32(x-1)(x-H_{2m})x^{n+1}W_{mn+j}^2 + 32(x-1)x^{n+1}W_{mn-m+j}^2 + 32(x-1)W_j^2 - 32(x-1)xW_{j-m}^2 + 2(W_1^2 + W_0^2 - 6W_1W_0)(x^n - 1)(H_{2m} - 2)x.$$

- (b) If $(1 + (-s)^{2m}x^2 - xH_{2m})((-s)^mx - 1) = u(x - a)(x - b)(x - c) = 0$ for some $u, a, b, c \in \mathbb{C}$ with $u \neq 0$ and $a \neq b \neq c$, i.e., $x = a$ or $x = b$ or $x = c$, then

$$\sum_{k=0}^n x^k W_{mk+j}^2 = \frac{\Omega_2}{32(3x^2 - 2(H_{2m} + 1)x + H_{2m} + 1)},$$

where

$$\Omega_2 = 32((x - H_{2m})x^{n+1} + (x - 1)((n + 2)x - (n + 1)H_{2m})x^n W_{mn+j}^2 + 32((n + 2)x - (n + 1))x^n W_{mn-m+j}^2 + 32W_j^2 - 32(2x - 1)W_{j-m}^2 + 2(W_1^2 + W_0^2 - 6W_1W_0)(x^n(n + 1) - 1)(H_{2m} - 2).$$

- (c) If $(1 + (-s)^{2m}x^2 - xH_{2m})((-s)^mx - 1) = u(x - a)^2(x - c) = 0$ for some $u, a, c \in \mathbb{C}$ with $u \neq 0$, $a \neq c$, when $x = c$, then

$$\sum_{k=0}^n x^k W_{mk+j}^2 = \frac{\Omega_3}{32(3x^2 - 2(H_{2m} + 1)x + H_{2m} + 1)},$$

where

$$\Omega_3 = 32((x - H_{2m})x^{n+1} + (x - 1)((n + 2)x - (n + 1)H_{2m})x^n W_{mn+j}^2 + 32((n + 2)x - (n + 1))x^n W_{mn-m+j}^2 + 32W_j^2 - 32(2x - 1)W_{j-m}^2 + 2(W_1^2 + W_0^2 - 6W_1W_0)(x^n(n + 1) - 1)(H_{2m} - 2),$$

and when $x = a$, then

$$\sum_{k=0}^n x^k W_{mk+j}^2 = \frac{\Omega_4}{64(3x - 1 - H_{2m})},$$

where

$$\Omega_4 = 32((n + 3)(n + 2)x^2 - x(n + 2)(n + 1)(H_{2m} + 1) + n(n + 1)H_{2m})x^{n-1}W_{mn+j}^2 + 32(n + 1)((2 + n)x^n - nx^{n-1})W_{mn-m+j}^2 - 64W_{j-m}^2 + 2n(n + 1)(W_1^2 + W_0^2 - 6W_1W_0)(H_{2m} - 2)x^{n-1}.$$

- (d) If $(1 + (-s)^{2m}x^2 - xH_{2m})((-s)^mx - 1) = u(x - a)^3 = 0$ for some $u, a \in \mathbb{C}$ with $u \neq 0$, i.e., $x = a$, then

$$\sum_{k=0}^n x^k W_{mk+j}^2 = \frac{\Omega_5}{192},$$

where

$$\Omega_5 = 32(n + 1)((n + 3)(n + 2)x^2 - n(n + 2)(H_{2m} + 1)x + n(n - 1)H_{2m})x^{n-2}W_{mn+j}^2 + 32n(n + 1)((n + 2)x + 1 - n)x^{n-2}W_{mn-m+j}^2 + 2(n - 1)n(n + 1)(H_{2m} - 2)(W_1^2 + W_0^2 - 6W_1W_0)x^{n-2}.$$

Proof. Take $r = 6$, $s = -1$ in Soykan [28], Theorem 2.1.. □

Note that (2.1) can be written in the following form

$$\sum_{k=1}^n x^k W_{mk+j}^2 = \frac{\Omega_6}{32(1 + x^2 - xH_{2m})(x - 1)},$$

where

$$\Omega_6 = 32(x - 1)(x - H_{2m})x^{n+1}W_{mn+j}^2 + 32((-s)^mx - 1)x^{n+1}W_{mn-m+j}^2 - 32(x - 1)(x - H_{2m})x W_j^2 - 32(x - 1)W_{j-m}^2 x + 2(W_1^2 + W_0^2 - 6W_1W_0)(x^n - 1)(H_{2m} - 2)x.$$

As special cases of m and j in the last theorem, one obtains the following proposition.

Proposition 2.2. For generalized balancing numbers (the case $r = 6, s = -1$) one has the following sum formulas for $n \geq 0$:

(a) ($m = 1, j = 0$)

If $(x - 1)(x^2 - 34x + 1) \neq 0$, i.e., $x \neq 1, x \neq 17 - 12\sqrt{2}, x \neq 17 + 12\sqrt{2}$, then

$$\sum_{k=0}^n x^k W_k^2 = \frac{\Delta}{(x-1)(x^2-34x+1)},$$

where

$$\Delta = (x-1)(x-34)x^{n+1}W_n^2 + (x-1)x^{n+1}W_{n-1}^2 + (x-1)W_0^2 - (x-1)x(W_1 - 6W_0)^2 + 2(W_1^2 + W_0^2 - 6W_1W_0)(x^n - 1)x,$$

and

if $(x-1)(x^2 - 34x + 1) = 0$, i.e., $x = 1$ or $x = 17 - 12\sqrt{2}$ or $x = 17 + 12\sqrt{2}$, then

$$\sum_{k=0}^n x^k W_k^2 = \frac{\Psi}{(3x^2 - 70x + 35)},$$

where

$$\Psi = ((x-34)x^{n+1} + (x-1)((n+2)x - 34(n+1))x^n W_n^2 + ((n+2)x - (n+1))x^n W_{n-1}^2 + W_0^2 - (2x-1)(W_1 - 6W_0)^2 + 2(W_1^2 + W_0^2 - 6W_1W_0)(x^n(n+1) - 1)$$

(b) ($m = 2, j = 0$)

If $(x-1)(x^2 - 1154x + 1) \neq 0$, i.e., $x \neq 1, x \neq 577 - 408\sqrt{2}, x \neq 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k W_{2k}^2 = \frac{\Delta}{(x-1)(x^2-1154x+1)},$$

where

$$\Delta = (x-1)(x-1154)x^{n+1}W_{2n}^2 + (x-1)x^{n+1}W_{2n-2}^2 + (x-1)W_0^2 - (x-1)x(6W_1 - 35W_0)^2 + 72(W_1^2 + W_0^2 - 6W_1W_0)(x^n - 1)x,$$

and

if $(x-1)(x^2 - 1154x + 1) = 0$, i.e., $x = 1$ or $x = 577 - 408\sqrt{2}$ or $x = 577 + 408\sqrt{2}$ then

$$\sum_{k=0}^n x^k W_{2k}^2 = \frac{\Psi}{3(x^2 - 770x + 385)},$$

where

$$\Psi = ((x-1154)x^{n+1} + (x-1)((n+2)x - 1154(n+1))x^n W_{2n}^2 + ((n+2)x - (n+1))x^n W_{2n-2}^2 + W_0^2 - (2x-1)(6W_1 - 35W_0)^2 + 72(W_1^2 + W_0^2 - 6W_1W_0)(x^n(n+1) - 1)$$

(c) ($m = 2, j = 1$)

If $(x-1)(x^2 - 1154x + 1) \neq 0$, i.e., $x \neq 1, x \neq 577 - 408\sqrt{2}, x \neq 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k W_{2k+1}^2 = \frac{\Delta}{(x-1)(x^2-1154x+1)},$$

where

$$\Delta = (x-1)(x-1154)x^{n+1}W_{2n+1}^2 + (x-1)x^{n+1}W_{2n-1}^2 + (x-1)W_1^2 - (x-1)x(W_1-6W_0)^2 + 72(W_1^2 + W_0^2 - 6W_1W_0)(x^n-1)x,$$

and

if $(x-1)(x^2-1154x+1) = 0$, i.e., $x = 1$ or $x = 577 - 408\sqrt{2}$ or $x = 577 + 408\sqrt{2}$ then

$$\sum_{k=0}^n x^k W_{2k+1}^2 = \frac{\Psi}{3(x^2-770x+385)},$$

where

$$\Psi = ((x-1154)x^{n+1} + (x-1)((n+2)x-1154(n+1))x^n)W_{2n+1}^2 + ((n+2)x-(n+1))x^nW_{2n-1}^2 + W_1^2 - (2x-1)(W_1-6W_0)^2 + 72(W_1^2 + W_0^2 - 6W_1W_0)(x^n(n+1)-1)$$

(d) $(m = -1, j = 0)$

If $(x-1)(x^2-34x+1) \neq 0$, i.e., $x \neq 1$, $x \neq 17 - 12\sqrt{2}$, $x \neq 17 + 12\sqrt{2}$, then

$$\sum_{k=0}^n x^k W_{-k}^2 = \frac{\Delta}{(x-1)(x^2-34x+1)},$$

where

$$\Delta = (x-1)x^{n+1}W_{-n+1}^2 + (x-1)(x-34)x^{n+1}W_{-n}^2 + (x-1)W_0^2 - (x-1)xW_1^2 + 2(W_1^2 + W_0^2 - 6W_1W_0)(x^n-1)x,$$

and

if $(x-1)(x^2-34x+1) = 0$, i.e., $x = 1$ or $x = 17 - 12\sqrt{2}$ or $x = 17 + 12\sqrt{2}$, then

$$\sum_{k=0}^n x^k W_{-k}^2 = \frac{\Psi}{(3x^2-70x+35)},$$

where

$$\Psi = ((n+2)x-(n+1))x^nW_{-n+1}^2 + ((x-34)x^{n+1} + (x-1)((n+2)x-34(n+1))x^n)W_{-n}^2 + W_0^2 - (2x-1)W_1^2 + 2(W_1^2 + W_0^2 - 6W_1W_0)(x^n(n+1)-1)$$

(e) $(m = -2, j = 0)$

If $(x-1)(x^2-1154x+1) \neq 0$, i.e., $x \neq 1$, $x \neq 577 - 408\sqrt{2}$, $x \neq 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k W_{-2k}^2 = \frac{\Delta}{(x-1)(x^2-1154x+1)},$$

where

$$\Delta = (x-1)x^{n+1}W_{-2n+2}^2 + (x-1)(x-1154)x^{n+1}W_{-2n}^2 + (x-1)W_0^2 - (x-1)x(W_0-6W_1)^2 + 72(W_1^2 + W_0^2 - 6W_1W_0)(x^n-1)x,$$

and

if $(x-1)(x^2-1154x+1) = 0$, i.e., $x = 1$ or $x = 577 - 408\sqrt{2}$ or $x = 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k W_{-2k}^2 = \frac{\Psi}{3(x^2-770x+385)},$$

where

$$\Psi = ((n+2)x-(n+1))x^nW_{-2n+2}^2 + ((x-1154)x^{n+1} + (x-1)((n+2)x-1154(n+1))x^n)W_{-2n}^2 + W_0^2 - (2x-1)(W_0-6W_1)^2 + 72(W_1^2 + W_0^2 - 6W_1W_0)(x^n(n+1)-1)$$

(f) ($m = -2, j = 1$)

If $(x - 1)(x^2 - 1154x + 1) \neq 0$, i.e., $x \neq 1, x \neq 577 - 408\sqrt{2}, x \neq 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k W_{-2k+1}^2 = \frac{\Delta}{(x-1)(x^2 - 1154x + 1)},$$

where

$$\Delta = (x-1)x^{n+1}W_{-2n+3}^2 + (x-1)(x-1154)x^{n+1}W_{-2n+1}^2 + (x-1)W_1^2 - (x-1)x(6W_0 - 35W_1)^2 + 72(W_1^2 + W_0^2 - 6W_1W_0)(x^n - 1)x,$$

and

if $(x-1)(x^2 - 1154x + 1) = 0$, i.e., $x = 1$ or $x = 577 - 408\sqrt{2}$ or $x = 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k W_{-2k+1}^2 = \frac{\Psi}{3(x^2 - 770x + 385)},$$

where

$$\Psi = ((n+2)x - (n+1))x^n W_{-2n+3}^2 + ((x-1154)x^{n+1} + (x-1)((n+2)x - 1154(n+1))x^n)W_{-2n+1}^2 + W_1^2 - (2x-1)(6W_0 - 35W_1)^2 + 72(W_1^2 + W_0^2 - 6W_1W_0)(x^n(n+1) - 1)$$

From the above proposition, one has the following corollary, which gives sum formulas of balancing numbers (take $W_n = B_n$ with $B_0 = 0, B_1 = 1$).

Corollary 2.3. For $n \geq 0$, balancing numbers have the following properties:

(a) ($m = 1, j = 0$)

If $(x-1)(x^2 - 34x + 1) \neq 0$, i.e., $x \neq 1, x \neq 17 - 12\sqrt{2}, x \neq 17 + 12\sqrt{2}$, then

$$\sum_{k=0}^n x^k B_k^2 = \frac{(x-1)(x-34)x^{n+1}B_n^2 + (x-1)x^{n+1}B_{n-1}^2 + x(2x^n - x - 1)}{(x-1)(x^2 - 34x + 1)},$$

and

if $(x-1)(x^2 - 34x + 1) = 0$, i.e., $x = 1$ or $x = 17 - 12\sqrt{2}$ or $x = 17 + 12\sqrt{2}$, then

$$\sum_{k=0}^n x^k B_k^2 = \frac{\Theta_1}{(3x^2 - 70x + 35)},$$

where

$$\Theta_1 = ((x-34)x^{n+1} + (x-1)((n+2)x - 34(n+1))x^n)B_n^2 + ((n+2)x - (n+1))x^n B_{n-1}^2 + 2(n+1)x^n - 2x - 1.$$

(b) ($m = 2, j = 0$)

If $(x-1)(x^2 - 1154x + 1) \neq 0$, i.e., $x \neq 1, x \neq 577 - 408\sqrt{2}, x \neq 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k B_{2k}^2 = \frac{(x-1)(x-1154)x^{n+1}B_{2n}^2 + (x-1)x^{n+1}B_{2n-2}^2 - 36x(-2x^n + x + 1)}{(x-1)(x^2 - 1154x + 1)}$$

and

if $(x-1)(x^2-1154x+1)=0$, i.e., $x=1$ or $x=577-408\sqrt{2}$ or $x=577+408\sqrt{2}$, then

$$\sum_{k=0}^n x^k B_{2k}^2 = \frac{\Theta_2}{3(x^2-770x+385)},$$

where

$$\Theta_2 = ((x-1154)x^{n+1} + (x-1)((n+2)x-1154(n+1))x^n)B_{2n}^2 + ((n+2)x-(n+1))x^n B_{2n-2}^2 + 36(2(n+1)x^n - 2x - 1).$$

(c) ($m=2, j=1$)

If $(x-1)(x^2-1154x+1) \neq 0$, i.e., $x \neq 1, x \neq 577-408\sqrt{2}, x \neq 577+408\sqrt{2}$, then

$$\sum_{k=0}^n x^k B_{2k+1}^2 = \frac{(x-1)(x-1154)x^{n+1}B_{2n+1}^2 + (x-1)x^{n+1}B_{2n-1}^2 - (-72x^{n+1} + x^2 + 70x + 1)}{(x-1)(x^2-1154x+1)},$$

and

if $(x-1)(x^2-1154x+1)=0$, i.e., $x=1$ or $x=577-408\sqrt{2}$ or $x=577+408\sqrt{2}$, then

$$\sum_{k=0}^n x^k B_{2k+1}^2 = \frac{\Theta_3}{3(x^2-770x+385)},$$

where

$$\Theta_3 = ((x-1154)x^{n+1} + (x-1)((n+2)x-1154(n+1))x^n)B_{2n+1}^2 + ((n+2)x-(n+1))x^n B_{2n-1}^2 + 2(36(n+1)x^n - x - 35).$$

(d) ($m=-1, j=0$)

If $(x-1)(x^2-34x+1) \neq 0$, i.e., $x \neq 1, x \neq 17-12\sqrt{2}, x \neq 17+12\sqrt{2}$, then

$$\sum_{k=0}^n x^k B_{-k}^2 = \frac{(x-1)x^{n+1}B_{-n+1}^2 + (x-1)(x-34)x^{n+1}B_{-n}^2 + x(2x^n - x - 1)}{(x-1)(x^2-34x+1)},$$

and

if $(x-1)(x^2-34x+1)=0$, i.e., $x=1$ or $x=17-12\sqrt{2}$ or $x=17+12\sqrt{2}$, then

$$\sum_{k=0}^n x^k B_{-k}^2 = \frac{\Theta_4}{(3x^2-70x+35)}$$

where

$$\Theta_4 = ((n+2)x-(n+1))x^n B_{-n+1}^2 + ((x-34)x^{n+1} + (x-1)((n+2)x-34(n+1))x^n)B_{-n}^2 + 2(n+1)x^n - 2x - 1.$$

(e) ($m=-2, j=0$)

If $(x-1)(x^2-1154x+1) \neq 0$, i.e., $x \neq 1, x \neq 577-408\sqrt{2}, x \neq 577+408\sqrt{2}$, then

$$\sum_{k=0}^n x^k B_{-2k}^2 = \frac{(x-1)x^{n+1}B_{-2n+2}^2 + (x-1)(x-1154)x^{n+1}B_{-2n}^2 - 36x(-2x^n + x + 1)}{(x-1)(x^2-1154x+1)}$$

and

if $(x-1)(x^2-1154x+1)=0$, i.e., $x=1$ or $x=577-408\sqrt{2}$ or $x=577+408\sqrt{2}$, then

$$\sum_{k=0}^n x^k B_{-2k}^2 = \frac{\Theta_5}{3(x^2-770x+385)},$$

where

$$\Theta_5 = ((n+2)x - (n+1))x^n B_{-2n+2}^2 + ((x-1154)x^{n+1} + (x-1)((n+2)x - 1154(n+1))x^n)B_{-2n}^2 + 36(2(n+1)x^n - 2x - 1).$$

(f) $(m=-2, j=1)$

If $(x-1)(x^2-1154x+1) \neq 0$, i.e., $x \neq 1$, $x \neq 577-408\sqrt{2}$, $x \neq 577+408\sqrt{2}$, then

$$\sum_{k=0}^n x^k B_{-2k+1}^2 = \frac{(x-1)x^{n+1}B_{-2n+3}^2 + (x-1)(x-1154)x^{n+1}B_{-2n+1}^2 + (72x^{n+1} - 1225x^2 + 1154x - 1)}{(x-1)(x^2-1154x+1)},$$

and

if $(x-1)(x^2-1154x+1)=0$, i.e., $x=1$ or $x=577-408\sqrt{2}$ or $x=577+408\sqrt{2}$, then

$$\sum_{k=0}^n x^k B_{-2k+1}^2 = \frac{\Theta_6}{3(x^2-770x+385)},$$

where

$$\Theta_6 = ((n+2)x - (n+1))x^n B_{-2n+3}^2 + ((x-1154)x^{n+1} + (x-1)((n+2)x - 1154(n+1))x^n)B_{-2n+1}^2 + 2(36(n+1)x^n - 1225x + 577).$$

Taking $W_n = H_n$ with $H_0 = 2, H_1 = 6$ in the last proposition, one has the following corollary, which presents sum formulas of modified Lucas-balancing numbers.

Corollary 2.4. For $n \geq 0$, modified Lucas-balancing numbers have the following properties:

(a) $(m=1, j=0)$

If $(x-1)(x^2-34x+1) \neq 0$, i.e., $x \neq 1$, $x \neq 17-12\sqrt{2}$, $x \neq 17+12\sqrt{2}$, then

$$\sum_{k=0}^n x^k H_k^2 = \frac{(x-1)(x-34)x^{n+1}H_n^2 + (x-1)x^{n+1}H_{n-1}^2 - 4(16x^{n+1} + 9x^2 - 26x + 1)}{(x-1)(x^2-34x+1)},$$

and

if $(x-1)(x^2-34x+1)=0$, i.e., $x=1$ or $x=17-12\sqrt{2}$ or $x=17+12\sqrt{2}$, then

$$\sum_{k=0}^n x^k H_k^2 = \frac{\Theta_7}{(3x^2-70x+35)},$$

where

$$\Theta_7 = ((x-34)x^{n+1} + (x-1)((n+2)x - 34(n+1))x^n)H_n^2 + ((n+2)x - (n+1))x^n H_{n-1}^2 - 8(8(n+1)x^n + 9x - 13).$$

(b) ($m = 2, j = 0$)

If $(x - 1)(x^2 - 1154x + 1) \neq 0$, i.e., $x \neq 1, x \neq 577 - 408\sqrt{2}, x \neq 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k H_{2k}^2 = \frac{(x-1)(x-1154)x^{n+1}H_{2n}^2 + (x-1)x^{n+1}H_{2n-2}^2 - 4(576x^{n+1} + 289x^2 - 866x + 1)}{(x-1)(x^2 - 1154x + 1)},$$

and

if $(x - 1)(x^2 - 1154x + 1) = 0$, i.e., $x = 1$ or $x = 577 - 408\sqrt{2}$ or $x = 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k H_{2k}^2 = \frac{\Theta_8}{3(x^2 - 770x + 385)},$$

where

$$\Theta_8 = ((x - 1154)x^{n+1} + (x - 1)((n + 2)x - 1154(n + 1))x^n)H_{2n}^2 + ((n + 2)x - (n + 1))x^n H_{2n-2}^2 - 8(288(n + 1)x^n + 289x - 433).$$

(c) ($m = 2, j = 1$)

If $(x - 1)(x^2 - 1154x + 1) \neq 0$, i.e., $x \neq 1, x \neq 577 - 408\sqrt{2}, x \neq 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k H_{2k+1}^2 = \frac{(x-1)(x-1154)x^{n+1}H_{2n+1}^2 + (x-1)x^{n+1}H_{2n-1}^2 - 36(64x^{n+1} + x^2 - 66x + 1)}{(x-1)(x^2 - 1154x + 1)},$$

and

if $(x - 1)(x^2 - 1154x + 1) = 0$, i.e., $x = 1$ or $x = 577 - 408\sqrt{2}$ or $x = 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k H_{2k+1}^2 = \frac{\Theta_9}{3(x^2 - 770x + 385)},$$

where

$$\Theta_9 = ((x - 1154)x^{n+1} + (x - 1)((n + 2)x - 1154(n + 1))x^n)H_{2n+1}^2 + ((n + 2)x - (n + 1))x^n H_{2n-1}^2 - 72(32(n + 1)x^n + x - 33).$$

(d) ($m = -1, j = 0$)

If $(x - 1)(x^2 - 34x + 1) \neq 0$, i.e., $x \neq 1, x \neq 17 - 12\sqrt{2}, x \neq 17 + 12\sqrt{2}$, then

$$\sum_{k=0}^n x^k H_{-k}^2 = \frac{(x-1)x^{n+1}H_{-n+1}^2 + (x-1)(x-34)x^{n+1}H_{-n}^2 - 4(16x^{n+1} + 9x^2 - 26x + 1)}{(x-1)(x^2 - 34x + 1)},$$

and

if $(x - 1)(x^2 - 34x + 1) = 0$, i.e., $x = 1$ or $x = 17 - 12\sqrt{2}$ or $x = 17 + 12\sqrt{2}$ then

$$\sum_{k=0}^n x^k H_{-k}^2 = \frac{\Theta_{10}}{(3x^2 - 70x + 35)},$$

where

$$\Theta_{10} = ((n + 2)x - (n + 1))x^n H_{-n+1}^2 + ((x - 34)x^{n+1} + (x - 1)((n + 2)x - 34(n + 1))x^n)H_{-n}^2 - 8(8(n + 1)x^n + 9x - 13).$$

(e) ($m = -2, j = 0$)

If $(x - 1)(x^2 - 1154x + 1) \neq 0$, i.e., $x \neq 1, x \neq 577 - 408\sqrt{2}, x \neq 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k H_{-2k}^2 = \frac{(x-1)x^{n+1}H_{-2n+2}^2 + (x-1)(x-1154)x^{n+1}H_{-2n}^2 - 4(576x^{n+1} + 289x^2 - 866x + 1)}{(x-1)(x^2 - 1154x + 1)},$$

and

if $(x - 1)(x^2 - 1154x + 1) = 0$, i.e., $x = 1$ or $x = 577 - 408\sqrt{2}$ or $x = 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k H_{-2k}^2 = \frac{\Theta_{11}}{3(x^2 - 770x + 385)},$$

where

$$\Theta_{11} = ((n+2)x - (n+1))x^n H_{-2n+2}^2 + ((x-1154)x^{n+1} + (x-1)((n+2)x - 1154(n+1))x^n)H_{-2n}^2 - 8(288(n+1)x^n + 289x - 433).$$

(f) ($m = -2, j = 1$)

If $(x - 1)(x^2 - 1154x + 1) \neq 0$, i.e., $x \neq 1, x \neq 577 - 408\sqrt{2}, x \neq 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k H_{-2k+1}^2 = \frac{(x-1)x^{n+1}H_{-2n+3}^2 + (x-1)(x-1154)x^{n+1}H_{-2n+1}^2 - 36(64x^{n+1} + 1089x^2 - 1154x + 1)}{(x-1)(x^2 - 1154x + 1)}$$

and

if $(x - 1)(x^2 - 1154x + 1) = 0$, i.e., $x = 1$ or $x = 577 - 408\sqrt{2}$ or $x = 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k H_{-2k+1}^2 = \frac{\Theta_{12}}{3(x^2 - 770x + 385)},$$

where

$$\Theta_{12} = ((n+2)x - (n+1))x^n H_{-2n+3}^2 + ((x-1154)x^{n+1} + (x-1)((n+2)x - 1154(n+1))x^n)H_{-2n+1}^2 - 72(32(n+1)x^n + 1089x - 577).$$

From the above proposition, one has the following corollary, which gives sum formulas of Lucas-balancing numbers (take $W_n = C_n$ with $C_0 = 1, C_1 = 3$).

Corollary 2.5. For $n \geq 0$, Lucas-balancing numbers have the following properties:

(a) ($m = 1, j = 0$)

If $(x - 1)(x^2 - 34x + 1) \neq 0$, i.e., $x \neq 1, x \neq 17 - 12\sqrt{2}, x \neq 17 + 12\sqrt{2}$, then

$$\sum_{k=0}^n x^k C_k^2 = \frac{(x-1)(x-34)x^{n+1}C_n^2 + (x-1)x^{n+1}C_{n-1}^2 - (16x^{n+1} + 9x^2 - 26x + 1)}{(x-1)(x^2 - 34x + 1)},$$

and

if $(x - 1)(x^2 - 34x + 1) = 0$, i.e., $x = 1$ or $x = 17 - 12\sqrt{2}$ or $x = 17 + 12\sqrt{2}$, then

$$\sum_{k=0}^n x^k C_k^2 = \frac{\Theta_{13}}{(3x^2 - 70x + 35)},$$

where

$$\Theta_{13} = ((x-34)x^{n+1} + (x-1)((n+2)x - 34(n+1))x^n)C_n^2 + ((n+2)x - (n+1))x^n C_{n-1}^2 - 2(8(n+1)x^n + 9x - 13).$$

(b) ($m = 2, j = 0$)

If $(x - 1)(x^2 - 1154x + 1) \neq 0$, i.e., $x \neq 1, x \neq 577 - 408\sqrt{2}, x \neq 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k C_{2k}^2 = \frac{(x-1)(x-1154)x^{n+1}C_{2n}^2 + (x-1)x^{n+1}C_{2n-2}^2 - (576x^{n+1} + 289x^2 - 866x + 1)}{(x-1)(x^2 - 1154x + 1)},$$

and

if $(x - 1)(x^2 - 1154x + 1) = 0$, i.e., $x = 1$ or $x = 577 - 408\sqrt{2}$ or $x = 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k C_{2k}^2 = \frac{\Theta_{14}}{3(x^2 - 770x + 385)},$$

where

$$\Theta_{14} = ((x - 1154)x^{n+1} + (x - 1)((n + 2)x - 1154(n + 1))x^n)C_{2n}^2 + ((n + 2)x - (n + 1))x^n C_{2n-2}^2 - 2(288(n + 1)x^n + 289x - 433).$$

(c) ($m = 2, j = 1$)

If $(x - 1)(x^2 - 1154x + 1) \neq 0$, i.e., $x \neq 1, x \neq 577 - 408\sqrt{2}, x \neq 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k C_{2k+1}^2 = \frac{(x-1)(x-1154)x^{n+1}C_{2n+1}^2 + (x-1)x^{n+1}C_{2n-1}^2 - 9(64x^{n+1} + x^2 - 66x + 1)}{(x-1)(x^2 - 1154x + 1)},$$

and

if $(x - 1)(x^2 - 1154x + 1) = 0$, i.e., $x = 1$ or $x = 577 - 408\sqrt{2}$ or $x = 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k C_{2k+1}^2 = \frac{\Theta_{15}}{3(x^2 - 770x + 385)},$$

where

$$\Theta_{15} = ((x - 1154)x^{n+1} + (x - 1)((n + 2)x - 1154(n + 1))x^n)C_{2n+1}^2 + ((n + 2)x - (n + 1))x^n C_{2n-1}^2 - 18(32(n + 1)x^n + x - 33).$$

(d) ($m = -1, j = 0$)

If $(x - 1)(x^2 - 34x + 1) \neq 0$, i.e., $x \neq 1, x \neq 17 - 12\sqrt{2}, x \neq 17 + 12\sqrt{2}$, then

$$\sum_{k=0}^n x^k C_{-k}^2 = \frac{(x-1)x^{n+1}C_{-n+1}^2 + (x-1)(x-34)x^{n+1}C_{-n}^2 - (16x^{n+1} + 9x^2 - 26x + 1)}{(x-1)(x^2 - 34x + 1)},$$

and

if $(x - 1)(x^2 - 34x + 1) = 0$, i.e., $x = 1$ or $x = 17 - 12\sqrt{2}$ or $x = 17 + 12\sqrt{2}$, then

$$\sum_{k=0}^n x^k C_{-k}^2 = \frac{\Theta_{16}}{(3x^2 - 70x + 35)},$$

where

$$\Theta_{16} = ((n + 2)x - (n + 1))x^n C_{-n+1}^2 + ((x - 34)x^{n+1} + (x - 1)((n + 2)x - 34(n + 1))x^n)C_{-n}^2 - 2(8(n + 1)x^n + 9x - 13).$$

(e) ($m = -2, j = 0$)

If $(x - 1)(x^2 - 1154x + 1) \neq 0$, i.e., $x \neq 1, x \neq 577 - 408\sqrt{2}, x \neq 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k C_{-2k}^2 = \frac{(x-1)x^{n+1}C_{-2n+2}^2 + (x-1)(x-1154)x^{n+1}C_{-2n}^2 - (576x^{n+1} + 289x^2 - 866x + 1)}{(x-1)(x^2 - 1154x + 1)},$$

and

if $(x - 1)(x^2 - 1154x + 1) = 0$, i.e., $x = 1$ or $x = 577 - 408\sqrt{2}$ or $x = 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k C_{-2k}^2 = \frac{\Theta_{17}}{3(x^2 - 770x + 385)},$$

where

$$\Theta_{17} = ((n+2)x - (n+1))x^n C_{-2n+2}^2 + ((x-1154)x^{n+1} + (x-1)((n+2)x - 1154(n+1))x^n)C_{-2n}^2 - 2(288(n+1)x^n + 289x - 433).$$

(f) ($m = -2, j = 1$)

If $(x - 1)(x^2 - 1154x + 1) \neq 0$, i.e., $x \neq 1, x \neq 577 - 408\sqrt{2}, x \neq 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k C_{-2k+1}^2 = \frac{(x-1)x^{n+1}C_{-2n+3}^2 + (x-1)(x-1154)x^{n+1}C_{-2n+1}^2 - 9(64x^{n+1} + 1089x^2 - 1154x + 1)}{(x-1)(x^2 - 1154x + 1)},$$

and

if $(x - 1)(x^2 - 1154x + 1) = 0$, i.e., $x = 1$ or $x = 577 - 408\sqrt{2}$ or $x = 577 + 408\sqrt{2}$, then

$$\sum_{k=0}^n x^k C_{-2k+1}^2 = \frac{\Theta_{18}}{3(x^2 - 770x + 385)},$$

where

$$\Theta_{18} = ((n+2)x - (n+1))x^n C_{-2n+3}^2 + ((x-1154)x^{n+1} + (x-1)((n+2)x - 1154(n+1))x^n)C_{-2n+1}^2 - 18(32(n+1)x^n + 1089x - 577).$$

Taking $x = 1$ in the last two corollaries, one gets the following corollary.

Corollary 2.6. For $n \geq 0$, balancing numbers, modified Lucas-balancing and Lucas-balancing numbers have the following properties:

1.

- (a) $\sum_{k=0}^n B_k^2 = \frac{1}{32}(33B_n^2 - B_{n-1}^2 - 2n + 1)$.
- (b) $\sum_{k=0}^n B_{2k}^2 = \frac{1}{1152}(1153B_{2n}^2 - B_{2n-2}^2 - 72n + 36)$.
- (c) $\sum_{k=0}^n B_{2k+1}^2 = \frac{1}{1152}(1153B_{2n+1}^2 - B_{2n-1}^2 - 72n)$.
- (d) $\sum_{k=0}^n B_{-k}^2 = \frac{1}{32}(-B_{-n+1}^2 + 33B_{-n}^2 - 2n + 1)$.
- (e) $\sum_{k=0}^n B_{-2k}^2 = \frac{1}{1152}(-B_{-2n+2}^2 + 1153B_{-2n}^2 - 72n + 36)$.
- (f) $\sum_{k=0}^n B_{-2k+1}^2 = \frac{1}{1152}(-B_{-2n+3}^2 + 1153B_{-2n+1}^2 - 72n + 1224)$.

2.

- (a) $\sum_{k=0}^n H_k^2 = \frac{1}{32}(33H_n^2 - H_{n-1}^2 + 64n + 32).$
- (b) $\sum_{k=0}^n H_{2k}^2 = \frac{1}{1152}(1153H_{2n}^2 - H_{2n-2}^2 + 2304n + 1152).$
- (c) $\sum_{k=0}^n H_{2k+1}^2 = \frac{1}{1152}(1153H_{2n+1}^2 - H_{2n-1}^2 + 2304n).$
- (d) $\sum_{k=0}^n H_{-k}^2 = \frac{1}{32}(-H_{-n+1}^2 + 33H_{-n}^2 + 64n + 32).$
- (e) $\sum_{k=0}^n H_{-2k}^2 = \frac{1}{1152}(-H_{-2n+2}^2 + 1153H_{-2n}^2 + 2304n + 1152).$
- (f) $\sum_{k=0}^n H_{-2k+1}^2 = \frac{1}{1152}(-H_{-2n+3}^2 + 1153H_{-2n+1}^2 + 2304n + 39168).$

3.

- (a) $\sum_{k=0}^n C_k^2 = \frac{1}{32}(33C_n^2 - C_{n-1}^2 + 16n + 8).$
- (b) $\sum_{k=0}^n C_{2k}^2 = \frac{1}{1152}(1153C_{2n}^2 - C_{2n-2}^2 + 576n + 288).$
- (c) $\sum_{k=0}^n C_{2k+1}^2 = \frac{1}{1152}(1153C_{2n+1}^2 - C_{2n-1}^2 + 576n).$
- (d) $\sum_{k=0}^n C_{-k}^2 = \frac{1}{32}(-C_{-n+1}^2 + 33C_{-n}^2 + 16n + 8).$
- (e) $\sum_{k=0}^n C_{-2k}^2 = \frac{1}{1152}(-C_{-2n+2}^2 + 1153C_{-2n}^2 + 576n + 288).$
- (f) $\sum_{k=0}^n C_{-2k+1}^2 = \frac{1}{1152}(-C_{-2n+3}^2 + 1153C_{-2n+1}^2 + 576n + 9792).$

3 Conclusion

Recently, there have been so many studies of the sequences of numbers in the literature. The sequences of numbers were widely used in many research areas, such as architecture, nature, art, physics, and engineering. In this work, sum identities were proved. The method used in this paper can be used for the other linear recurrence sequences, too. We have written sum identities in terms of the generalized balancing sequence. Then, we have presented the formulas as special cases, the corresponding identity for the balancing, modified Lucas-balancing, and Lucas-balancing numbers. All the listed identities in the corollaries may be proved by induction, but that proof method gives no clue about their discovery. We have provided proofs to show how these identities were discovered in general.

Computations of the Frobenius norm, spectral norm, maximum column length norm, and maximum row length norm of circulant (r-circulant, geometric circulant, semicirculant) matrices with the generalized m -step Fibonacci sequences require the sum of the numbers of the sequences. So, our results can be used to study r-circulant matrices with m -order linear recurrence sequences.

Acknowledgements

No funding is available for this research.

Conflict of Interest

The authors have no conflicts of interest to declare.

References

- [1] A. BEHERA AND G. K. PANDA, *On the Square Roots of Triangular Numbers*, Fibonacci Quart. **37**(2) (1999), 98–105.
- [2] A. BERCZES, K. LIPTAI AND I. PINK, *On Generalized Balancing Sequences*, Fibonacci Quart. **48**(2) (2010), 121–128. .
- [3] P. CATARINO, H. CAMPOS AND P. VASCO, *On Some Identities for Balancing and Cobalancing Numbers*, Ann. Math. Inform. **45** (2015), 11–24.
- [4] R.K. DAVALA AND G.K. PANDA, *On Sum and Ratio Formulas for Balancing Numbers*, J. Indian Math. Soc. (N.S.) **82**(1-2) (2015), 23–32.
- [5] A.F. HORADAM, *A Generalized Fibonacci Sequence*, Amer. Math. Monthly **68** (1961), 455–459.
- [6] A.F. HORADAM, *Basic Properties of a Certain Generalized Sequence of Numbers*, Fibonacci Quart. **3**(3) (1965), 161–176.
- [7] A.F. HORADAM, *Special Properties of The Sequence $w_n(a, b; p, q)$* , Fibonacci Quart. **5**(5) (1967), 424–434.
- [8] A.F. HORADAM, *Generating functions for powers of a certain generalized sequence of numbers*, Duke Math. J. **32** (1965), 437–446.
- [9] R. FRONTCAK, *Identities for Generalized balancing Numbers*, Notes Number Theory Discrete Math. **25**(2) (2019) 169–180. DOI:10.7546/nntdm.2019.25.2.169-180
- [10] R. KESKIN AND O. KARAATLI, *Some New Properties of Balancing Numbers and Square Triangular Numbers*, J. Integer Seq. **15** (2012), 1–13.
- [11] T. KOVACS, K. LIPTAI AND P. OLAJOS, *On $(a; b)$ -balancing Numbers*, Publ. Math. Debrecen **77**(3-4) (2010), 485–498.
- [12] K. LIPTAI, *Fibonacci Balancing Numbers*, Fibonacci Quart. **42**(4) (2004), 330–340.
- [13] K. LIPTAI, F. LUCA, A. PINTER AND L. SZALAY, *Generalized Balancing Numbers*, Indag. Math. (N.S.) **20**(1) (2009), 87–100.
- [14] P. OLAJOS, *Properties of Balancing, Cobalancing and Generalized Balancing Numbers*, Ann. Math. Inform. **37** (2010), 125–138.
- [15] A. ÖZKOÇ, *Tridiagonal Matrices via k -Balancing Number*, British J. Math. Comput. Sci. **10**(4) (2015), 1–11.
- [16] G.K. PANDA, *Some Fascinating Properties of Balancing Numbers*, In Proc. of Eleventh Internat. Conference on Fibonacci Numbers and Their Applications, Congr. Numer. **194** (2009), 185–189.
- [17] G.K. PANDA, T. KOMATSU AND R.K. DAVALA, *Reciprocal Sums of Sequences Involving Balancing and Lucas-Balancing Numbers*, Math. Rep. **20**(70) (2018), 201–214.
- [18] P.K. RAY, *On the Properties of Lucas-balancing Numbers by Matrix Method*, Sigmae **3**(1) (2014), 1–6.

- [19] P.K. RAY, *Balancing Sequences of Matrices with Application to Algebra of balancing Numbers*, Notes Number Theory Discrete Math. **20**(1) (2014), 49–58.
- [20] P.K. RAY, *Some Congruences for Balancing and Lucas-Balancing Numbers and Their Applications*, Integers **14** (2014), 1–8.
- [21] P.K. RAY, *Balancing and Lucas-Balancing Sums by Matrix Methods*, Math. Rep. **17**(2) (2015), 225–233.
- [22] P.K. RAY, S. PATEL AND M.K. MANDAL, *Identities for Balancing Numbers Using Generating Function and Some New Congruence Relations*, Notes Number Theory Discrete Math. **22**(4) (2016), 41–48.
- [23] P.K. RAY AND J. SAHU, *Generating Functions for Certain Balancing and Lucas-Balancing Numbers*, Palestine J. Math. **5**(2) (2016), 122–129.
- [24] P.K. RAY, *Balancing Polynomials and Their Derivatives*, Ukr. Math. J. **69** (2017), 646–663.
- [25] S.G. RAYAGURU AND G.K. PANDA, *Sum Formulas Involving Powers of Balancing and Lucas-balancing Numbers*, J. Ind. Math. Soc. **86**(1-2) (2019), 137–160.
- [26] S.G. RAYAGURU AND G.K. PANDA, *Sum Formulas Involving Powers of Balancing and Lucas-balancing Numbers-II*, Notes Number Theory Discrete Math. **25**(3) (2019), 102–110. DOI: [10.7546/nntdm.2019.25.3.102-110](https://doi.org/10.7546/nntdm.2019.25.3.102-110)
- [27] N.J.A. SLOANE, *The On-Line Encyclopedia of Integer Sequences*, Published electronically at <http://oeis.org>.
- [28] Y. SOYKAN, *Generalized Fibonacci Numbers with Indices in Arithmetic Progression and Sum of Their Squares: the Sum Formula $\sum_{k=0}^n x^k W_{mk+j}^2$* , J. Adv. Math. Comput. Sci. **36**(6) (2021), 30–62. DOI: [10.9734/JAMCS/2021/v36i630371](https://doi.org/10.9734/JAMCS/2021/v36i630371)
- [29] Y. SOYKAN, *A Study on Generalized Balancing Numbers*, Asian J. Adv. Res. Rep. **15**(5) (2021), 78–100. DOI: [10.9734/AJARR/2021/v15i530401](https://doi.org/10.9734/AJARR/2021/v15i530401)